

Online Research @ Cardiff

This is an Open Access document downloaded from ORCA, Cardiff University's institutional repository: <https://orca.cardiff.ac.uk/id/eprint/76873/>

This is the author's version of a work that was submitted to / accepted for publication.

Citation for final published version:

Bidon, Tobias, Schreck, Nancy, Hailer, Frank ORCID: <https://orcid.org/0000-0002-2340-1726>, Nilsson, Maria A. and Janke, Axel 2015. Genome-wide search identifies 1.9 Mb from the polar bear Y chromosome for evolutionary analyses. *Genome Biology and Evolution* 7 (7) , pp. 2010-2022. 10.1093/gbe/evv103 file

Publishers page: <http://dx.doi.org/10.1093/gbe/evv103>
<<http://dx.doi.org/10.1093/gbe/evv103>>

Please note:

Changes made as a result of publishing processes such as copy-editing, formatting and page numbers may not be reflected in this version. For the definitive version of this publication, please refer to the published source. You are advised to consult the publisher's version if you wish to cite this paper.

This version is being made available in accordance with publisher policies.

See

<http://orca.cf.ac.uk/policies.html> for usage policies. Copyright and moral rights for publications made available in ORCA are retained by the copyright holders.



Genome-Wide Search Identifies 1.9 Mb from the Polar Bear Y Chromosome for Evolutionary Analyses

Tobias Bidon^{1,2,†}, Nancy Schreck^{1,†}, Frank Hailer^{1,3}, Maria A. Nilsson¹, and Axel Janke^{1,4,*}

¹Senckenberg Biodiversity and Climate Research Centre Frankfurt, Frankfurt am Main, Germany

²International Graduate School of Science and Engineering (IGSSE), Technische Universität München, Garching, Germany

³School of Biosciences, Cardiff University, Wales, United Kingdom

⁴Institute for Ecology, Evolution & Diversity, Goethe University Frankfurt, Germany

*Corresponding author: E-mail: ajanke@senckenberg.de.

[†]These authors contributed equally to this work.

Accepted: May 23, 2015

Abstract

The male-inherited Y chromosome is the major haploid fraction of the mammalian genome, rendering Y-linked sequences an indispensable resource for evolutionary research. However, despite recent large-scale genome sequencing approaches, only a handful of Y chromosome sequences have been characterized to date, mainly in model organisms. Using polar bear (*Ursus maritimus*) genomes, we compare two different *in silico* approaches to identify Y-linked sequences: 1) Similarity to known Y-linked genes and 2) difference in the average read depth of autosomal versus sex chromosomal scaffolds. Specifically, we mapped available genomic sequencing short reads from a male and a female polar bear against the reference genome and identify 112 Y-chromosomal scaffolds with a combined length of 1.9 Mb. We verified the *in silico* findings for the longer polar bear scaffolds by male-specific *in vitro* amplification, demonstrating the reliability of the average read depth approach. The obtained Y chromosome sequences contain protein-coding sequences, single nucleotide polymorphisms, microsatellites, and transposable elements that are useful for evolutionary studies. A high-resolution phylogeny of the polar bear patriline shows two highly divergent Y chromosome lineages, obtained from analysis of the identified Y scaffolds in 12 previously published male polar bear genomes. Moreover, we find evidence of gene conversion among *ZFX* and *ZFY* sequences in the giant panda lineage and in the ancestor of ursine and tremarctine bears. Thus, the identification of Y-linked scaffold sequences from unordered genome sequences yields valuable data to infer phylogenomic and population-genomic patterns in bears.

Key words: sex chromosome, patriline, male inheritance, Ursidae, Y chromosome, divergence.

Introduction

Genomic sequence data have become an important resource for evolutionary biology, and new sequenced genomes are becoming available at increasing speed. The mammalian genome consists of autosomes, sex chromosomes and mitochondrial (mt) DNA, which are differentially inherited. These parts of the genome can thus provide information about distinctive aspects of a species' evolutionary history (Chesser and Baker 1996; Veeramah and Hammer 2014).

For technical reasons, the maternally inherited mtDNA has been a standard tool to study evolutionary processes in model and nonmodel organisms (Wilson et al. 1985). Consequently, the first available genomic resources for evolutionary studies were fully sequenced mt genomes (Anderson et al. 1981;

Janke et al. 1994). The paternally inherited counterpart of mtDNA is the male-specific Y chromosome, one of two sex chromosomes in the mammalian genome. Similar to mtDNA, the Y chromosome is haploid, lacks interchromosomal recombination for most of its length, and is uniparentally inherited. These properties allow the inference of long and high-resolution haplotypes, enabling researchers to trace the evolutionary history of male lineages over time (Jobling and Tyler-Smith 2003; Wei et al. 2013). Biparentally inherited autosomes provide the largest amount of sequence data, but their phylogenetic analysis can be complicated by reticulate evolution (Posada et al. 2002).

Polar bears have recently been in the focus of genome scale evolutionary analyses and genomic sequences have been used

to address the evolution, population history, and unique adaptations of this high arctic mammal (Hailer et al. 2012; Miller et al. 2012; Cahill et al. 2013, 2015; Cronin et al. 2014; Liu et al. 2014). Furthermore, short Y-chromosomal sequences and six Y-linked microsatellites in polar and brown bears were used to investigate the distribution of male-specific genetic variation across their ranges (Bidon et al. 2014). The application of Y-chromosomal markers is particularly interesting in many mammals, because sex-specific differences in dispersal behavior are predicted to affect phylogeographic and population genetic conclusions that have so far been drawn almost exclusively from female-inherited mtDNA.

Y-chromosomal sequences are important in studies of evolutionary history, chromosome structure, and forensic applications (Jobling and Tyler-Smith 2003; Kayser 2007; Hallast et al. 2013). The Y chromosome's unique evolutionary viewpoint has been used to investigate patterns of domestication and migration, for example, in horses and dogs (Sacks et al. 2013; Wallner et al. 2013), and to study human phylogeography and migration (Wei et al. 2013; Scozzari et al. 2014; Van Oven et al. 2014).

Despite the wealth of genomic data, identification of large amounts of Y-chromosomal sequences from high throughput sequencing data is rarely done. Genome sequences are usually ordered into scaffolds, without information about their relative orientation or chromosomal origin, because thoroughly annotated reference genomes and physical maps are still lacking for most taxonomic groups. In addition, many mammalian genomes have been sequenced from female individuals, to obtain equal coverage of autosomes and the X chromosome (Hughes and Rozen 2012), but also for technical difficulties relating to the assembly and the high amount of repetitive and ampliconic sequences on Y chromosomes (Willard 2003; Bachtrog 2013). This has hampered sequencing, assembly, identification, and application of Y-chromosomal markers (Greminger et al. 2010). As a consequence, complete Y chromosome sequences are only published for four mammalian species: Human, chimpanzee, rhesus macaque, and mouse (Skaletsky et al. 2003; Hughes et al. 2010, 2012; Soh et al. 2014). This list is complemented by large-scale analyses of Y-chromosomal sequences for dog, cat, marmoset, rat, bull, opossum, *Drosophila*, and medaka fish (Kondo et al. 2006; Carvalho et al. 2009; Li et al. 2013; Bellott et al. 2014).

In this study, we utilize a previously published polar bear reference assembly that is based on a male sequenced at high coverage (Li et al. 2011), and available short sequence reads from additional male and female polar bears (Miller et al. 2012). We identify Y-linked scaffolds by 1) searching for sequence similarity using known mammalian Y-linked gene sequences and 2) identifying scaffolds with sex-specific sequencing coverage characteristics indicative of Y linkage. The latter approach makes use of differences in the expected sequence coverage of male versus female sequence reads on autosomal, X-chromosomal, and Y-chromosomal reference

scaffolds. We apply stringent quality filters to minimize false positives, that is, scaffolds wrongly identified as Y-linked. In addition, in vitro amplification of the longest candidate scaffolds confirmed the in silico findings. We demonstrate that genome scale Y-chromosomal sequences can be reliably identified from high-throughput sequencing data, also in organisms lacking a chromosome-based physical map of the genome.

Materials and Methods

We used two different approaches to identify Y-chromosomal sequences in the recently published polar bear genome assembly from a male individual that was sequenced at 101-fold coverage (Li et al. 2011). This assembly has a size of 2.3 Gb and is arranged into 72,214 scaffolds with an N50 value of 15.9 Mb. Information regarding chromosomal locations and the relative orientation of the scaffolds is not available. Thus, it is unknown which of the scaffolds are of Y-chromosomal origin. In the following, we refer to this genome assembly as the "polar bear assembly," and to the scaffolds of this assembly by their respective scaffold ID numbers.

Similarity Search of Y-Linked Genes Lists Candidate Scaffolds

The first approach was to use 32 genes known to be Y-linked in other mammals as queries for a similarity search in the polar bear assembly. Exon sequences from human (*Homo sapiens*), mouse (*Mus musculus*), chimpanzee (*Pan troglodytes*), and dog (*Canis lupus familiaris*) were downloaded from GenBank for these genes (supplementary table S1, Supplementary Material online). Similarity between exon sequences and the scaffolds of the polar bear assembly was identified using Basic Local Alignment Search Tool (BLAST), analyzing one exon at a time.

Scaffolds from the polar bear assembly were extracted from the list of BLAST hits according to the following criteria: 1) The scaffold with the lowest *E* value (expect value) for a particular exon relative to all other scaffolds in the list and 2) scaffolds with $\geq 95\%$ sequence similarity compared with the scaffold with the lowest *E* value, with the additional constraint that the difference in alignment length of exon and scaffold (compared with the scaffold with the lowest *E* value) must not exceed 5%. We then obtained the exact position of each exon on its respective scaffold by realigning exon and scaffold using ClustalW. Only scaffolds with a sequence identity of $\geq 80\%$ between scaffold and exon were kept (Table 1).

In addition, in vitro validated male-specific polar bear sequences from five known Y-linked genes (exons and introns, *AMELY*, *KDM5D* [*SMCY*], *SRY*, *UBA1Y*, *ZFY*; supplementary table S1, Supplementary Material online) (Nakagome et al. 2008; Pagès et al. 2008, 2009) were downloaded and used

Table 1

Polar Bear Scaffolds Showing Similarity to 18 Mammalian Y-Linked Genes

Nr.	Scaffold (ID)	Scaffold Size (kb)	Gene
1	13	26,707	<i>RPS4Y</i>
2	20	22,125	<i>EIF1AY, EIF2S3Y, USP9Y, ZFY</i>
3	46	15,941	<i>TBL1Y</i>
4	53	14,458	<i>SLY</i>
5	104	6,801	<i>NLGN4Y, PRKY, TBL1Y</i>
6	105	6,717	<i>RBMY1A1</i>
7	115	5,608	<i>AMELY</i>
8	134	4,673	<i>UBA1Y, UTY</i>
9	184	2,589	<i>DDX3Y, USP9Y</i>
10	186	2,578	<i>PCDH11Y</i>
11	253	821	<i>RPS4Y, RPS4Y2</i>
12	297 ^a	391	<i>EIF1AY, KDM5D</i>
13	301	351	<i>KDM5D</i>
14	309 ^a	317	<i>DDX3Y, USP9Y, UTY,</i>
15	318 ^a (3836) ^b	237	<i>EIF2S3Y, KDM5D, USP9Y, ZFY^b</i>
16	369 ^a	104	<i>RBMY1A1</i>
17	389 ^a	77	<i>AMELY</i>
18	403 ^a	63	<i>UBA1Y</i>
19	579 ^a	21	<i>SRY</i>
20	605 ^a	19	<i>UBA1Y</i>
21	646 ^a	15	<i>EIF2S3Y</i>
22	4889	0.9	<i>UBA1Y</i>
23	6612	0.7	<i>AMELY</i>

^aScaffolds have an AD-ratio indicative of Y-linkage and were validated in vitro to be male-specific.

^bThe entire length of scaffold 3836 (1 kb; with similarity to *ZFY*) is included within scaffold 318 with 100% identity.

as a query against the polar bear assembly using BLAT, with default parameters. Polar bear sequences from the two X-linked genes *ZFX* and *AMELX* (Pagès et al. 2009) were used to differentiate between Y and X gametologs, that is, homologous gene copies on the X and the Y chromosomes (supplementary table S1, Supplementary Material online).

Average Depth Ratio for Identification of Y-Linked, X-Linked, and Autosomal Scaffolds

In a second approach to identify Y-linked scaffolds, we utilized previously published short sequence reads from whole-genome sequence data of one female (SRX155950/PB06) and one male (SRX155954/PB10) polar bear. The two polar bear individuals had been sequenced at similar sequence depth (~12×) on an Illumina HiSeq 2000 platform, generating paired-end reads (101 bp) with an insert size of about 400 bp (Miller et al. 2012). The AD ratio approach is based on differences in the relative numbers of X and Y chromosomes between females (2-0) and males (1-1), whereas both sexes carry two copies of each autosome. As unique Y-chromosomal sequences are not present in a female genome, reads obtained from genome sequencing of female and male individuals

should map with characteristic sex-specific patterns to scaffolds from the Y chromosome, the X chromosome, and the autosomes. The expected differences in sequencing coverage were utilized primarily to identify Y-chromosomal scaffolds in the polar bear assembly, but our approach also allowed the assignment of anonymous scaffolds from the polar bear assembly as autosomal or X-linked.

Short read sequences were evaluated for residual adapter sequences and low-quality bases were clipped off the read-ends using FastQC v 0.10.0 (Andrews 2010) and sickle (Joshi and Fass 2011). BWA (Li and Durbin 2009) was used for the reference-guided mapping of the cleaned reads against the polar bear assembly. Using Samtools (Li et al. 2009), we merged read data from separate sequencing runs of the same sample into one single BAM file per individual. Picard (<http://picard.sourceforge.net/>, last accessed February 9, 2015) was used to mark duplicated reads, and realignment of reads was performed in GATK v2.3 (McKenna et al. 2010).

After mapping, the mpileup module of samtools was used to calculate the read depth at each position on a given scaffold for the male and the female genomes. Scaffolds without mapped reads or with low mapping quality ($n=614$), and scaffolds that were shorter than 1 kb ($n=68,017$; ~15 Mb) were disregarded and not considered in the downstream analyses. For the remaining scaffolds (≥ 1 kb, $n=3,583$), the average read depth was calculated: We determined the sum of the depth values at ambiguity-free scaffold positions (no "N") with ≤ 50 reads per position, and divided this by the number of ambiguity free scaffold positions.

Finally, the AD-ratio of each scaffold was calculated by dividing the average read depth in the female individual by the average read depth in the male individual (1). A normalization factor adjusted the number of female and male reads to each other (2): To this end, we divided the total number of reads (quality ≥ 20) in the female BAM file by that of the male BAM file.

For each given scaffold, average sequencing depth for the female and male genome was calculated using the following formulas:

$$\text{AD-ratio} = \frac{\text{average-depth}_{\text{female}}}{(\text{average-depth}_{\text{male}} \times \text{norm})} \quad (1)$$

$$\text{norm} = \frac{\text{total number of reads}_{\text{female}}}{\text{total number of reads}_{\text{male}}} \quad (2)$$

The normalization factor is used to enable comparison of read depth of individual scaffolds among individuals, despite possible differences in genome-wide sequencing coverage between them. Using this normalization factor, the male and female genomes are standardized to the same genome-wide average coverage. The AD-ratio is zero for perfectly mapped Y chromosome scaffolds, one for autosomal and two for X-linked scaffolds. For graphical representation, we

Table 2

Y-Chromosomal Scaffolds ≥ 10 kb Identified by the AD-Ratio

Nr.	Scaffold ID	Size (kb)	AD-Ratio	Similarity to Y-Linked Gene
1	297	391	0.284	<i>EIF1AY, KDM5D</i>
2	309	317	0	<i>DDX3Y, USP9Y, UTY</i>
3	318	237	0.285	<i>EIF2S3Y, KDM5D, USP9Y, ZFY</i>
4	322	217	0.252	—
5	369	104	0.18	<i>RBMY1A1</i>
6	389	77	0.16	<i>AMELY</i>
7	393	70	0.12	—
8	403	63	0	<i>UBA1Y</i>
9	420	54	0	—
10	519	31	0.198	—
11	579	21	0.075	<i>SRY</i>
12	596	20	0	—
13	605	19	0	<i>UBA1Y</i>
14	613	18	0.158	—
15	632	16	0.205	—
16	646	15	0	<i>EIF2S3Y</i>
17	657	14	0	—
18	771	10	0.135	—
19	795	10	0	—
20	813	10	0	—

NOTE.—The male-specificity of all scaffolds listed here has been validated in vitro. Additional scaffolds (<10 kb) are shown in [supplementary table S2, Supplementary Material](#) online.

combined scaffolds with different AD-ratios into bins of size 0.02.

In Vitro Validation of Putative Y-Linked Scaffolds in Different Bear Species

To verify the male-specificity of scaffolds identified by the in silico analysis, we PCR-amplified fragments from 20 Y-scaffolds (table 2) in at least one male and one female individual of each of three closely related ursine bears: Polar bear, brown bear, and American black bear (*U. americanus*). In addition, amplification of fragments from two X-linked and two autosomal scaffolds as identified by the AD-ratio approach was made in both male and female bears to verify their non-Y-chromosomal origin. Before amplification, newly designed primers ([supplementary table S7, Supplementary Material](#) online) were tested in silico for unique binding by aligning the forward and reverse sequences against the scaffolds of the polar bear assembly using BLASTn. Scaffolds were defined as being Y chromosome specific when one clear amplification product was detected in males, but no amplicons or only low-intensity bands of different sizes were observed in females. In vitro experiments included touchdown PCRs (see [supplementary material, Supplementary Material](#) online) and agarose gel-electrophoresis to verify the expected size of the amplicons. Each PCR setup contained a no-template control.

Repetitive Element Estimation in the Polar Bear Genome and on Y-Linked Scaffolds

The amount of transposable elements (TE) on 14 of the larger validated scaffolds (scaffold IDs: 297, 309, 318, 322, 369, 393, 389, 403, 420, 519, 579, 605, 646, and 657; 1.6 Mb) was identified using RepeatMasker (<http://www.repeatmasker.org>, last accessed February 9, 2015) using the carnivore library (Smit et al. 1996). RepeatMasker with the carnivore library was also used to identify microsatellites with a minimum of 15 repeat units ([supplementary table S6, Supplementary Material](#) online).

Analysis of X–Y Gene Conversion in Bears

The partial *ZFY* and *ZFX* exon sequences of all ursid species from Pagès et al. (2009) were downloaded from GenBank and aligned with homologous sequences from other mammals (397 bp) in Geneious 8.0.3 (Biomatters, Auckland, New Zealand). Based on the model suggested by jModeltest2 (Darriba et al. 2012), HKY+4 G, phylogenetic trees were constructed in Geneious, and a statistical parsimony network was generated in TCS (Clement et al. 2000).

Calculation of Polar Bear Patrilineal Phylogeny

Five Y-linked scaffolds (IDs: 309, 322, 389, 393, and 403) with a combined size of 743 kb were used to reconstruct the phylogenetic relationship of 12 polar bear individuals sampled in Svalbard (Norway) and Alaska. The Y-linked sequences were used to estimate the divergence time of the lineages within polar bears, using one American black and one brown bear as outgroup. Short reads of all 14 bear individuals (Miller et al. 2012) were retrieved from databases ([supplementary table S8, Supplementary Material](#) online) and mapped to the polar bear assembly as described above. The individuals have been labeled according to their respective description in the short read archive ([supplementary table S8, Supplementary Material](#) online). The five respective scaffolds together with previously mapped short reads were extracted using Samtools and loaded into Geneious 8.0.3 (Biomatters). Geneious was then used to create a consensus sequence for each individual, to align those consensus sequences, and to remove alignment columns containing ambiguous sites and gaps, respectively. Additionally, the alignments were manually inspected to find and remove columns where only one individual contained multiple differentiating sites adjacent to each other. This strict filtering reduced the size of the alignments by approximately 30% (see below).

A NeighborNet network was calculated in SplitsTree 4.12.6 (Huson and Bryant 2006) based on a 511-kb-long alignment of the concatenated Y-sequences of 12 polar bears. BEAST 2.1.3 (Bouckaert et al. 2014) was used to estimate divergence times among polar bears, using a strict clock model, a Yule tree model, and a uniform prior of 343–479 ka, based on the relatively young population divergence between brown and

polar bears (Liu et al. 2014). An additional calibration scenario employed a fixed mutation rate obtained from human Y chromosomes: 0.76×10^9 /site/year (Fu et al. 2014). We used the GTR+I substitution model as indicated by the Bayesian Information Criterion in jModeltest 2.1.1. Convergence was checked in Tracer (ESS > 200). The concatenated alignment comprising 506 kb included 12 polar bears, one brown bear, and one black bear. This alignment was thus slightly shorter than the polar bear alignment, due to ambiguous sites and gaps introduced by the inclusion of additional individuals/taxa.

Results

We identified a total of 1.9 Mb of Y-chromosomal sequence data in the polar bear assembly, located on 112 different scaffolds. The scaffolds were identified by applying two different approaches: 1) The search for similarity of known Y-linked genes, and 2) comparison of the AD-ratio of reads from male and female genomes.

The Similarity Search Identified 23 Putative Y-Chromosomal Scaffolds

The first approach identified scaffolds in the polar bear assembly that showed similarity to known Y-linked gene sequences from four different mammals (human, mouse, chimpanzee, and dog). Exons from 18 of 32 Y-linked candidate genes that were blasted against the polar bear assembly identified polar bear scaffold sequences above a threshold of 80% identity (table 1 and [supplementary table S1, Supplementary Material online](#)). The hits were distributed across 23 scaffolds, ranging from 0.7 to 26,707 kb in size (table 1 and fig. 1). The full sequence length of scaffold 3836 (1,069 bp) had an identical sequence stretch on scaffold 318 (237 kb), with no nucleotide mismatches. Thus, we do not report scaffold 3836 as a distinct scaffold, although it is a separate entry in the current polar bear assembly.

Six sequences of five Y-linked genes from polar bear Y chromosomes (Nakagome et al. 2008; Pagès et al. 2008, 2009) aligned to the polar bear assembly with 98.9–100% identity. A 227-bp fragment from *ZFY* and a 49-bp fragment from *KDM5D* (*SMCY*) were uncharacterized ("N") in the polar bear assembly.

We found that ten query genes had similarity to two or more different scaffolds in the polar bear assembly, thereby creating combinations of scaffolds that contain stretches of homologous sequence ([supplementary table S4, Supplementary Material online](#)). These scaffold combinations consisted of one in vitro validated Y-scaffold and one (or two) scaffold(s) with an AD-ratio expected for X-chromosomal linkage, indicating sequential homology between the Y-chromosomal and other scaffolds. For instance, the *ZFY* exon sequences mapped to both scaffold 318 and scaffold 20 with similar identity (99.2% vs. 99.5%). The gametologous polar bear *ZFX* sequence also mapped to both these scaffolds,

at the same location as *ZFY*. However, when using less conserved intronic sequences from polar bears ([supplementary table S1, Supplementary Material online](#)) in a BLAT search against the polar bear assembly, scaffolds 318 (containing *ZFY*, Y-linked) and scaffold 20 (containing *ZFX*, X-linked) were clearly diagnosable.

Phylogenetic analyses of *ZFX/ZFY* sequences in mammals showed that the X- and Y-linked copies of giant panda (*Ailuropoda melanoleuca*) form a cluster, and that all ursine and tremarctine *ZFX/ZFY* sequences form a second cluster of closely related sequences ([supplementary fig. S1, Supplementary Material online](#)). Ursid sequences thus clustered together, regardless of their X- or Y-chromosomal origin. Other mammals clustered outside the ursid variation.

Based on the similarity of known Y-linked candidate gene sequences from different mammals, the similarity search provided us with a list of 23 scaffolds that might potentially be located on the polar bear Y chromosome. However, 12 of these scaffolds were identified to be autosomal or X-linked, due to their respective AD-ratios (see below, fig. 1 and [supplementary table S4, Supplementary Material online](#)).

The Average-Depth Ratio Identified 112 Y-Chromosomal Scaffolds

Most scaffolds had an AD-ratio of either approximately 1 or approximately 2, indicative of autosomal and X-chromosomal scaffolds, respectively (fig. 2 and [supplementary fig. S2, Supplementary Material online](#)). The combined sequence length of all putative autosomal scaffolds ≥ 1 kb ($0.7 < \text{AD-ratio} < 1.3$; $n = 2,618$) was approximately 2.18 Gb, and putative X-linked scaffolds ($1.7 < \text{AD-ratio} < 2.3$; $n = 214$) amounted to approximately 109 Mb. At an AD-ratio of zero, which is the expected AD-ratio for Y-linked scaffolds, we detected 90 scaffolds with a combined sequence length of 686 kb (fig. 2 and [supplementary fig. S2, Supplementary Material online, table 2 and supplementary table S2, Supplementary Material online](#)). An additional 22 scaffolds with a combined sequence length of 1.21 Mb showed AD-ratios ≤ 0.3 , of which 11 were amplified in vitro, all showing male-specific amplification (table 2). Thus, applying a relaxed AD-ratio cutoff of ≤ 0.3 , thereby allowing for a certain proportion of wrongly mapped reads, identified 112 Y-linked scaffolds, comprising 1.9 Mb of Y-chromosomal sequence.

Nine scaffolds totaling 1.24 Mb were identified by both approaches (fig. 1 and [supplementary table S3, Supplementary Material online](#)). Among the scaffolds obtained exclusively from the similarity search, one had an AD-ratio of exactly zero, but it was less than 1 kb (scaffold ID 6612; 794 bp). Four putative Y-chromosomal scaffolds from the similarity search had an AD-ratio of approximately 1, indicating autosomal origin. Eight scaffolds from the similarity

search had an AD-ratio of approximately 2, indicating X-linked origin. For one scaffold identified by the similarity search, neither male nor female reads mapped with sufficient quality (scaffold ID 4889), precluding any linkage classification.

In Vitro Amplification Validates All Tested Y-Linked Scaffolds as Being Male-Specific

The male-specificity of the longest putative Y-linked scaffolds ($n = 20$) was additionally evaluated in vitro by PCR amplification (table 2). At least one fragment of 635–800 bp sequence length of each of the scaffolds was PCR-amplified using male DNA samples along with female DNA controls of each brown, polar and black bears. All 20 scaffolds showed male-specific PCR amplification, defined as the occurrence of a clear amplicon of a distinct size in males but not in females. In female DNA samples, the Y-chromosomal fragments could either not be amplified (scaffold IDs 309, 318, 322, 369, 389, 393, 420, 579, 596, 605, 613, 632, and 813) or the observed amplicons were smaller, with multiple low-intensity (unspecific) bands/smears on agarose gels (scaffold IDs 297, 403, 519, 646, 657, 771, and 795). For comparison, we validated two fragments with putative autosomal (scaffold IDs 236 and 267) or X-

linkage (scaffold IDs 301 and 253), based on results from the AD-ratio approach. Markers on these putatively non Y-linked scaffolds could be PCR-amplified in both male and female DNA samples, and showed clear amplicons of the same sizes in both sexes.

High Abundance of Repetitive Elements on the Y-Linked Scaffolds

Overall, TEs covered 54.38% of the total length of the 14 Y-scaffolds used in this analysis (supplementary tables S5 and S6, Supplementary Material online, fig. 3 and supplementary fig. S3, Supplementary Material online). The majority of the TE sequences represents placental mammalian LINE-1 (38%) or the carnivore CAN-SINEs (7.8%). The average LINE-1 coverage of the polar bear genome is 16.93%, thus LINE-1 covered nearly twice as much sequence on the Y chromosome scaffolds compared with the entire genome. In addition, one full-length LINE-1 copy, the L1-1_AME, with a length = 6,021 bp was found on scaffold 297 (fig. 3). The full-length L1-1_AME is likely to have been recently active, due to the presence of only two stop codons in the endonuclease/reverse transcriptase encoding ORF2. The abundance




FIG. 1.—Identified scaffolds in the polar bear assembly. (A) Scaffolds identified by the similarity search, the AD-ratio, and by both approaches (overlap). Scaffolds ≥ 10 kb are shown by their ID numbers. Details for 92 additionally Y-linked scaffolds (< 10 kb, combined length: ~ 170 kb) are listed in supplementary table S2, Supplementary Material online. Some scaffolds identified by the similarity search showed AD-ratio characteristic of autosomal linkage (red) or X-linkage (blue). Scaffolds with an asterisk (*) have been verified in vitro to be male-specific. Two asterisks indicate scaffolds that show PCR amplification in both sexes. No reads mapped with sufficient mapping quality to scaffold 4889, so its AD-ratio could not be calculated, and scaffold 6612 was shorter than 1 kb. (B) AD-ratios of X-linked (blue), autosomal (red), and Y-linked (green) scaffolds.




FIG. 2.—AD-ratio histogram of polar bear scaffolds. (A) Distribution of AD-ratios of scaffolds ≥ 1 kb, and their combined size shown in bins of width 0.02. Autosomal scaffolds cluster around an AD-ratio of 1 (red), X-linked scaffolds around 2 (blue). The stippled box highlights the region shown enlarged in (B). Enlargement of the box in (A). Scaffolds below the threshold of 0.3 are identified as Y-linked (green). Scaffolds unassigned to chromosomal classes are shown in black.




FIG. 3.—Annotations of Y-linked scaffolds 297 and 318. Exons homologous to mouse and human are shown in red. Previously published Y-linked polar bear sequences are shown in pink. The repeat unit of each microsatellite is indicated and regions with greater than 200 bp of consecutive “N” are highlighted in gray. Due to the high abundance, only placental mammalian non-LTR retrotransposons ≥ 500 bp (LINEs) and ≥ 100 bp (SINEs) were plotted. The maps of additional scaffolds are shown in [supplementary figure S3, Supplementary Material](#) online.

of repetitive regions along the Y chromosome, and the positions of homologous regions to candidate gene sequences are shown in figure 3, exemplary for two long Y-chromosomal scaffolds (scaffold IDs 297 and 318). Corresponding maps for 12 additional Y-scaffolds are provided in [supplementary figure S3, Supplementary Material](#) online. We found a higher abundance of LINE-1 elements and a lower abundance of older

LINE-2 and LINE-3 elements on the Y chromosome compared with the whole genome. Moreover, a higher abundance of carnivore-specific SINEs as compared with ancestral Mammalian Interspersed Repeats (MIR) was detected. We identified a similar amount of long terminal repeats (LTRs)/endogenous retroviruses (ERVs) on the Y chromosome and the whole genome, whereas less DNA transposons were

identified on the Y chromosome compared with the whole genome. We identified 115 microsatellites with at least 15 repeat units, many of which are likely to show intraspecific polymorphism and are thus useful for population genetic studies, covering 0.3% of the combined length of all Y-scaffolds (supplementary table S6, Supplementary Material online).

Phylogenetic Analyses Identify Two Distinct Male Polar Bear Lineages

Phylogenetic analysis of 511 kb Y chromosome sequence in 12 polar bears identified two highly divergent paternal lineages (fig. 4A), with two individuals (AK4 and PB16) being clearly separated from the remaining ten polar bears. This separation does not correspond to geography, as both major lineages occur in Alaska and Svalbard (Norway). Some individuals have a considerable number of unique substitutions (e.g., PB16: 55 substitutions) relative to 101 substitutions separating the two lineages. Our Bayesian analysis yielded a phylogenetic tree with high posterior support for all major

nodes (fig. 4B), showing two distinct patrilineal clades within polar bears. Based on the demographic split of brown and polar bears at 343–479 ka (Liu et al. 2014), we obtained a median divergence time estimate for the split of these two clades at 0.12 Ma (95% highest posterior density (HPD): 0.10–0.15). The split between brown and polar bears was estimated at 0.40 Ma (0.34–0.47), and the divergence of the black bear at 1.190 Ma (0.99–1.44). Using a fixed mutation rate as an alternative calibration scenario, older divergence time estimates were obtained: The split within polar bears was estimated at 0.22 Ma (0.19–0.25), the split between brown and polar bears at 0.70 Ma (0.65–0.76), and the divergence of the black bear 2.13 Ma (2.03–2.23).

Discussion

The Y chromosome is poorly characterized in most mammals, including the carnivoran bear family. We used the polar bear reference assembly to identify a large amount of Y-linked sequence, totaling 1.9 Mb distributed across 112 Y-linked




FIG. 4.—Phylogenomic analysis of approximately 0.5-Mb Y-chromosomal sequence from 12 polar bears. Geographic origins of the polar bear individuals are denoted by AK (Alaska) and PB (Svalbard). (A) NeighborNet analysis. (B) Time-calibrated Bayesian coalescent-based phylogeny from BEAST. Numbers at nodes indicate the median of the divergence time in million years ago, with 95% highest posterior density in brackets. Dots at nodes indicate posterior probability greater than 0.99. The scale axis is in units of million years ago. Note that approximately 80% older dates were retrieved from an alternative calibration scenario (see text).

scaffolds. We did so by applying a similarity search with mammalian Y-linked genes and by analyzing differences in sequencing coverage (AD-ratio) on Y-chromosomal, autosomal, and X-linked scaffolds. We verified these *in silico* results *in vitro* by validating male-specificity by PCR amplification for 20 of the largest scaffolds, corresponding to 1.7 Mb of Y-chromosomal sequence (fig. 1 and tables 1 and 2).

The AD-Ratio Approach Can Reliably Identify Scaffolds from Autosomes, the X and the Y Chromosomes

Using the AD-ratio approach, the majority of scaffolds in the polar bear assembly could be assigned to one of three chromosomal classes. We identified 2.18 Gb of autosomal sequence, which is close to the total size of the polar bear assembly of 2.3 Gb (fig. 2). The X-linked scaffolds amounted to 109 Mb, which approximates two-thirds the size of the human X chromosome (Ross et al. 2005). This scaffold class included the 12 scaffolds that were previously identified as being X-linked in bears (Cahill et al. 2013). The amount of identified autosomal and X-linked sequences thus fit the expectations for a typical mammalian genome of approximately 2–3 Gb (Rogers and Gibbs 2014). Only approximately 0.2% (~5.3 Mb) of the polar bear assembly remained unassigned, because AD-ratios for these scaffolds were beyond our thresholds for autosomal, X-chromosomal, and Y-chromosomal sequences. This illustrates the reliability of the AD-ratio approach, and its suitability to screen a genome assembly for the three chromosome classes. The 1.9 Mb identified to be Y-linked is a considerable amount of Y-chromosomal sequence, given the lack of Y-linked genomic sequences for many mammals, the generally small size of the mammalian Y chromosome, and its highly repetitive nature that impedes assembly. We likely underestimate the total amount of Y-linked sequences (see below), and 1.9 Mb represents only a small fraction of the entire polar bear Y chromosome. The size of Y chromosomes differs considerably among mammals and even among carnivores, but Y chromosomes are typically longer than 20 Mb (e.g., dog: 20 Mb, cat: 45 Mb; Li et al. 2013). Although the size of the polar bear Y chromosome has not yet been determined, it appears to be about half the physical size of the X chromosome in metaphase spreads (O'Brien et al. 2006).

The similarity search identified 23 scaffolds as being Y-linked; however, later inspection indicated that only nine of these had an AD ratio indicative of Y-linkage (fig. 1 and table 1 and supplementary table S3, Supplementary Material online). The AD-ratio approach yielded 112 Y-linked scaffolds and thus proved to be more efficient than the similarity search in terms of scaffold numbers. However, the nine scaffolds identified by both approaches total 1.24 Mb, which is more than 60% of the entire Y-linked sequence data. Although a similarity search is technically simple, successful, and easily applied, several drawbacks are associated with this approach.

We based our selection of query genes on their previous description as being Y-linked in other mammals, implicitly assuming the presence of these genes also on the polar bear Y chromosome. However, Y chromosomes can differ in their gene content across taxa, and lineage-specific sets of Y-linked genes exist (Murphy et al. 2006; Cortez et al. 2014). Indeed, we found that Y-linked genes that are absent in carnivores, for example, *NLGN4Y* (Cortez et al. 2014), were also absent in polar bears. In contrast, genes that are widespread throughout placental mammals, and occur in other carnivores (dog and cat; e.g., *ZFY*, *UTY*, *EIF1AY*) (Cortez et al. 2014), are those genes that are actually found on Y-linked scaffolds in the polar bear assembly (fig. 3 and supplementary fig. S3, Supplementary Material online). Currently limited knowledge of gene contents on the Y chromosomes of different mammalian lineages is therefore still restricting the efficiency of similarity-based approaches for the identification of Y-chromosomal scaffolds.

In several cases, the similarity search produced hits to more than one scaffold (supplementary table S4, Supplementary Material online). For example, a search with *ZFY* sequences yielded similarity to scaffold 318 (containing *ZFY*) and scaffold 20 (containing *ZFX*). The scaffolds in such groups all had AD-ratios characteristic of either Y- or X-linkage (supplementary table S4, Supplementary Material online). Most Y-linked genes on these scaffolds are classified as X-degenerate in humans (supplementary table S1, Supplementary Material online). These genes are relics of the ancient autosomes from which the mammalian X and Y chromosomes evolved, and are thus expected to show homology between the X and Y chromosomes (Skaletsky et al. 2003). In contrast, *RBMY* (scaffold IDs 369 and 105; supplementary table S4, Supplementary Material online) is classified as ampliconic in humans, and such genes normally lack X-linked counterparts. *RBMY*, however, is one of the two ampliconic genes with an X-linked homolog in humans (*RBMX*), explaining its detection on an X-linked scaffold. These findings illustrate the high degree of sequence similarity between some sex chromosome gametologs (homologous genes on the two sex chromosomes), and the common evolutionary history of Y and X chromosomes, deriving from an ancestral pair of autosomes. Moreover, four scaffolds with sequence similarity to Y-linked genes, but an AD-ratio indicative of autosomal (or pseudoautosomal) origin, were identified (supplementary table S3, Supplementary Material online). Interestingly, one of these genes (*RPS4Y*, on scaffold: 13) is in close proximity to the pseudoautosomal region (PAR) on the small arm of the human Y chromosome (Skaletsky et al. 2003). The location of the PAR is not known in polar bears, but genes in regions recombining with the X chromosome would hinder correct identification of Y-linked scaffolds by the AD-ratio approach.

The similarity search is further complicated by the high degree of similarity between some gametologous genes. This is exemplified by *ZFY/ZFX* genes, for which we were

initially not able to differentiate between the respective Y- and X-scaffolds based on exon sequences. The more rapidly evolving intron sequences, however, allowed us to differentiate between Y- and X-linked scaffolds (supplementary fig. S1, Supplementary Material online). With a more stringent set of candidate genes, that is, carnivore-specific Y-linked genes, the reliability of the similarity search can be improved, and the search for intronic sequences would allow for a better differentiation between gametologs on the two sex chromosomes. A drawback of a similarity search based on Y-linked gene sequences from other taxa is that scaffolds consisting of exclusively intergenic sequence cannot be identified. This is an important limitation of similarity search approaches, because mammalian Y chromosomes are generally gene poor, with only 78 protein-coding genes in humans (Bachtrog 2013). Indeed, 103 of 112 Y-linked scaffolds were solely identified by their AD-ratio. Nevertheless, four of five Y-linked scaffolds with a size of greater than 100 kb were also identified by the similarity search (fig. 1 and table 2). Assemblies with fewer but larger scaffolds will thus be more amenable to accurate detection of Y-linkage by a similarity search approach.

The Structure of the Y Chromosome Complicates Identification of Y-Linked Sequences

The heterochromatic, highly repetitive regions of a genome usually remain unassembled in whole-genome sequencing projects. Some Y chromosomes contain extended regions of largely uncharacterized heterochromatin, for example, human and *Drosophila* (Bachtrog 2013). Other Y chromosomes are largely euchromatic, for example, mouse and chimpanzee, but even the euchromatic regions are enriched for ampliconic sequences containing duplicated genes (Skaletsky et al. 2003; Hughes et al. 2010; Soh et al. 2014). Accurate sequence assembly is therefore inherently difficult for the Y chromosome, and sequence similarity to the X and possibly other chromosomes further complicates the identification of a distinct Y-linked sequence. Therefore, high-quality Y chromosome reference sequence assemblies are so far lacking from most mammalian genome sequencing projects.

The identification of Y-linked scaffolds has previously been achieved by in silico search for known Y-linked genes and massive in vitro PCR-based verification in *Drosophila* and *Anopheles* (Carvalho et al. 2000; Krzywinski et al. 2004). Moreover, Y-linked sequences can be retrieved by subtracting the scaffolds of the homogametic from the heterogametic assembly (Chen et al. 2014). Approaches based on some measure of the coverage depth of sequence reads on Y-linked scaffolds (Carvalho et al. 2003; Chen et al. 2012), for example, the “Y chromosome genome scan” (Carvalho and Clark 2013), or on the number of alignments in males and females, the “chromosome quotient” (Hall et al. 2013), have also been applied.

The occurrence of gene conversion, where a gene copy on one chromosome is overwritten by the information from the other chromosome, further complicates identification of chromosome-specific sequences. This process appears also to occur in the bear lineage (supplementary fig. S1, Supplementary Material online). Compared with the human and dog outgroups, the tremarctine and ursine ZFY and ZFX sequences cluster together, and not with human and dog ZFY and ZFX, respectively. Additionally, the ZFY and ZFX sequences from giant panda (*Ailuropoda melanoleuca*) are more closely related to each other than any gene copy is to those from ursine and tremarctine bears. A likely explanation for these observations is that gene conversion has occurred in the ancestral giant panda lineage as well as in the lineage leading to tremarctine and ursine bears. Considering the divergence times of ursid lineages, these two conversion events occurred in the Miocene, more than 12 and 6 Ma, respectively (Kutschera et al. 2014). The occurrence of gene conversion between sex chromosomes has been described in various mammalian lineages such as primates and felids (Slattery et al. 2000; Rosser et al. 2009; Trombetta et al. 2014), including ZFX/ZFY.

TEs on the sex chromosomes pose yet another challenge for accurate assembly and identification of chromosome-specific sequences. Mammalian genomes contain large amounts of TEs that propagate through different mechanisms. The human genome has over 44% of TEs (Lander et al. 2001), whereas the polar bear genome consists of 39.2% TEs (supplementary table S5, Supplementary Material online). Previous studies have shown that there is a preferential insertion of some TEs (primate-specific LINE1 and Alu elements) on the human and chimpanzee X and Y chromosomes (Kvikstad and Makova 2010). The same distribution is observed on polar bear Y chromosome scaffolds, as there is a high abundance of LINE-1 and the carnivore-specific Can-SINEs (Walters-Conte et al. 2011) compared with the autosomes (supplementary table S5, Supplementary Material online). The ancestral TEs, such as LINE-2, LINE-3 and MIR elements which were active before the split between marsupial and placental mammals (Smit and Riggs 1995), are found in very low numbers on the polar bear Y scaffolds. The ERV and DNA transposons seem to accumulate more evenly across the genome than LINE-1 and Can-SINEs, as there are only small differences between Y-chromosomal and autosomal scaffolds. The reason for the preferential accumulation on the sex chromosomes has been attributed to male and female germline TE integrations occurring before meiotic sex chromosome inactivation (Kvikstad and Makova 2010).

Due to the repetitive nature of the Y chromosome, assembly methods will likely produce numerous smaller scaffolds and collapse repetitive sequences into chimeric scaffolds that actually comprise multicopy sequences. Indeed, stretches of very high sequence coverage were found on many of the Y-linked scaffolds. Moreover, long and highly repetitive regions

of the Y chromosome might be entirely missing from the assembly. TEs and X-transposed sequences on the Y chromosome likely cause a proportion of female reads from polar bear X chromosome and autosomes to be falsely mapped to Y-linked scaffolds, due to the high similarity among such regions. This produces AD-ratios greater than zero for these true Y-scaffolds. Finally, due to the paucity of information on bear sex chromosomes, we cannot exclude the possibility of recent stratum formation, with the existence of segments that have not yet attained a high level of divergence between the Y and X chromosomes. Yet, recent stratum formation or added genes from autosomes have not been reported for the well-studied Y chromosomes of two other carnivores, cat and dog (Cortez et al. 2014). A strict AD-ratio threshold of exactly zero is therefore likely to produce many false negatives. Our employed relaxed AD-ratio threshold of ≤ 0.3 yielded an additional 22 scaffolds, 11 of them tested and verified in vitro (table 2) to be of Y-chromosomal origin.

Assembly artifacts resulting from the repetitive nature of the Y chromosome imply that we likely underestimate the actual number and length of the identified Y-linked sequences. The AD-ratio approach should thus not be seen as an attempt to identify all Y-linked sequences in bears, nor to determine the size of the polar bears' Y chromosome. Rather, the approach is an effective means to identify sequences that demonstrably have a high probability of being Y-linked and that can be used for evolutionary studies.

Y-Chromosomal Sequences Provide a High Resolution Patrilineal Perspective on Polar Bear Evolutionary History

Our phylogenetic analyses of Y-linked scaffold sequences provide a patrilineal view on polar bear evolution that support a previously identified pattern of two distinct Y-chromosomal lineages in polar bears, PO1.1 and PO2 (Bidon et al. 2014) (fig. 4). The large amount of analyzed sequence data provides high resolution of individual lineages, with many haplotype-specific substitutions. Our divergence time estimation places the split of these two polar bear clades around the Eemian interglacial period (0.12–0.13 Ma), implying that the two lineages separated long before the last glacial maximum (approximately 18–25 ka). The clear separation into two paternal lineages indicates an ancient population structuring in polar bears, possibly due to the separation into multiple refugia during glaciation cycles, similar to other arctic species (Flagstad and Røed 2003).

The divergence time was estimated using a recently published date on the population split between brown and polar bears (343–479 ka; Liu et al. 2014). This demographic split is expected to be younger than estimates based on the coalescence of allelic lineages, for example, the 338–934 ka estimated by Hailer et al. (2012). The lower effective population size of the Y chromosome implies that coalescence of Y-lineages occurs faster than that of autosomal lineages. Therefore,

the Y-chromosomal gene tree might track the demographic splits of the species more closely.

It is noteworthy that our divergence estimate of the black bear patriline (0.99–1.44 Ma 95% HPD) is relatively young in this calibration scenario. The fossil record suggests a first occurrence of the black bear lineage at least 1.8 Ma (Kurtén and Anderson 1980). In principle, a Y-specific mutation rate would be a reasonable alternative calibration method. However, an independent mutation rate for the ursid Y chromosome has not yet been determined, and lineage-specific rates in mammals make the adoption of a Y-specific rate from another taxon unreliable. Applying a recent estimate for the mutation rate of the human Y chromosome, we obtained even older divergence time estimates for the patrilines of polar bears (0.19–0.25 Ma), of brown and polar bears (0.65–0.76 Ma), and of the black bear lineage (2.03–2.23 Ma). These dates are broadly consistent with other estimates of genomic divergence times for these splits (Hailer et al. 2012; Cahill et al. 2013; Cronin et al. 2014), and more in line with the fossil record of American black bears.

Short Y-linked sequences were recently used as markers for sex determination in bears (Bidon et al. 2013), phylogeographic analyses of brown and polar bear brother lineages (Bidon et al. 2014), and phylogenetic analyses of all eight bear species (Kutschera et al. 2014). Sequences on Y-chromosomal scaffolds have thus already proven to be a reliable resource for studying the evolutionary history of polar bears and other members of the ursid family.

Conclusions

The analyses of Y-chromosomal scaffolds provided a high-resolution view on the patrilineal relationship within polar bears, identifying two highly distinct clades that separated during the middle Pleistocene. A preferential accumulation of younger TEs on the polar bear Y chromosome could be shown. As more and more genomes become available in the form of reference assemblies and short read archives, straightforward in silico strategies to identify sex-linked sequences from these data can now be applied in many species. Overall, the AD-ratio approach seems to be highly specific and preferable for a reliable identification of Y chromosome scaffolds. It can be used as long as a reference assembly of the heterogametic sex, and short reads of one male and one female are available.

Supplementary Material

Supplementary tables S1–S8 and figures S1–S3 are available at *Genome Biology and Evolution* online (<http://www.gbe.oxfordjournals.org/>).

Acknowledgments

This work was supported by LOEWE Landes-Offensive zur Entwicklung Wissenschaftlich-ökonomischer Exzellenz; the

Arthur und Aenne Feindt-Stiftung, Hamburg; and the Leibniz-Association. The authors thank Steven Fain, Nicolas Lecomte, and Frank Zachos for providing samples, and the anonymous reviewers for their constructive comments. Jón Baldur Hlíðberg kindly provided the polar bear painting.

Literature Cited

- Anderson S, et al. 1981. Sequence and organization of the human mitochondrial genome. *Nature* 290:457–465.
- Andrews S. 2010. FastQC: a quality control tool for high throughput sequence data.
- Bachtrog D. 2013. Y-chromosome evolution: emerging insights into processes of Y-chromosome degeneration. *Nat Rev Genet.* 14:113–124.
- Bellott DW, et al. 2014. Mammalian Y chromosomes retain widely expressed dosage-sensitive regulators. *Nature* 508:494–499.
- Bidon T, et al. 2013. A sensitive and specific multiplex PCR approach for sex identification of ursine and tremarctine bears suitable for non-invasive samples. *Mol Ecol Resour.* 13:362–368.
- Bidon T, et al. 2014. Brown and polar bear Y chromosomes reveal extensive male-biased gene flow within brother lineages. *Mol Biol Evol.* 31:1353–1363.
- Bouckaert R, et al. 2014. BEAST 2: a software platform for Bayesian evolutionary analysis. *PLoS Comput Biol.* 10:e1003537.
- Cahill JA, et al. 2013. Genomic evidence for island population conversion resolves conflicting theories of polar bear evolution. *PLoS Genet.* 9:e1003345.
- Cahill JA, et al. 2015. Genomic evidence of geographically widespread effect of gene flow from polar bears into brown bears. *Mol Ecol.* 24:1205–1217.
- Carvalho AB, Clark AG. 2013. Efficient identification of Y chromosome sequences in the human and *Drosophila* genomes. *Genome Res.* 23:1894–1907.
- Carvalho AB, et al. 2003. Y chromosome and other heterochromatic sequences of the *Drosophila melanogaster* genome: how far can we go? *Genetica* 117:227–237.
- Carvalho AB, Koerich LB, Clark AG. 2009. Origin and evolution of Y chromosomes: *Drosophila* tales. *Trends Genet.* 25:270–277.
- Carvalho AB, Lazzaro BP, Clark AG. 2000. Y chromosomal fertility factors kl-2 and kl-3 of *Drosophila melanogaster* encode dynein heavy chain polypeptides. *Proc Natl Acad Sci U S A.* 97:13239–13244.
- Chen N, Bellott DW, Page DC, Clark AG. 2012. Identification of avian W-linked contigs by short-read sequencing. *BMC Genomics* 13:183.
- Chen S, et al. 2014. Whole-genome sequence of a flatfish provides insights into ZW sex chromosome evolution and adaptation to a benthic lifestyle. *Nat Genet.* 46:253–260.
- Chesser RK, Baker RJ. 1996. Effective sizes and dynamics of uniparentally and diparentally inherited genes. *Genetics* 144:1225–1235.
- Clement M, Posada D, Crandall K. 2000. TCS: a computer program to estimate gene genealogies. *Mol Ecol.* 9:1657–1659.
- Cortez D, et al. 2014. Origins and functional evolution of Y chromosomes across mammals. *Nature* 508:488–493.
- Cronin MA, et al. 2014. Molecular phylogeny and SNP variation of polar bears (*Ursus maritimus*), brown bears (*U. arctos*), and black bears (*U. americanus*) derived from genome sequences. *J Hered.* 105:312–323.
- Darriba D, Taboada GL, Doallo R, Posada D. 2012. jModelTest 2: more models, new heuristics and parallel computing. *Nat Methods.* 9:772.
- Flagstad Ø, Røed KH. 2003. Refugial origins of reindeer (*Rangifer tarandus* L.) inferred from mitochondrial DNA sequences. *Evolution* 57:658–670.
- Fu Q, et al. 2014. Genome sequence of a 45,000-year-old modern human from western Siberia. *Nature* 514:445–449.
- Greminger MP, Krützen M, Schelling C, Pienkowska-Schelling A, Wandeler P. 2010. The quest for Y-chromosomal markers—methodological strategies for mammalian non-model organisms. *Mol Ecol Resour.* 10:409–420.
- Hailer F, et al. 2012. Nuclear genomic sequences reveal that polar bears are an old and distinct bear lineage. *Science* 336:344–347.
- Hall AB, et al. 2013. Six novel Y chromosome genes in *Anopheles* mosquitoes discovered by independently sequencing males and females. *BMC Genomics* 14:273.
- Hallast P, Balaesque P, Bowden GR, Ballereau S, Jobling M. 2013. Recombination dynamics of a human Y-chromosomal palindrome: rapid GC-biased gene conversion, multi-kilobase conversion tracts, and rare inversions. *PLoS Genet.* 9:e1003666.
- Hughes JF, et al. 2010. Chimpanzee and human Y chromosomes are remarkably divergent in structure and gene content. *Nature* 463:536–539.
- Hughes JF, et al. 2012. Strict evolutionary conservation followed rapid gene loss on human and rhesus Y chromosomes. *Nature* 483:82–86.
- Hughes JF, Rozen S. 2012. Genomics and genetics of human and primate Y chromosomes. *Annu Rev Genomics Hum Genet.* 13:83–108.
- Huson DH, Bryant D. 2006. Application of phylogenetic networks in evolutionary studies. *Mol Biol Evol.* 23:254–267.
- Janke A, Feldmaier-Fuchs G, Thomas WK, von Haeseler A, Pääbo S. 1994. The marsupial mitochondrial genome and the evolution of placental mammals. *Genetics* 137:243–256.
- Jobling M, Tyler-Smith C. 2003. The human Y chromosome: an evolutionary marker comes of age. *Nat Rev Genet.* 4:598–612.
- Joshi NA, Fass JN. 2011. Sickle: a sliding-window, adaptive, quality-based trimming tool for FastQ files. Available from: <https://github.com/najoshi/sickle>
- Kayser M. 2007. Uni-parental markers in human identity testing including forensic DNA analysis. *Biotechniques* 43:Sxv–Sxxi.
- Kondo M, et al. 2006. Genomic organization of the sex-determining and adjacent regions of the sex chromosomes of medaka. *Genome Res.* 16:815–826.
- Krzywinski J, Nusskern DR, Kern MK, Besansky NJ. 2004. Isolation and characterization of Y chromosome sequences from the African malaria mosquito *Anopheles gambiae*. *Genetics* 166:1291–1302.
- Kurtén B, Anderson E. 1980. Pleistocene mammals of North America. New York: Columbia University Press.
- Kutschera VE, et al. 2014. Bears in a forest of gene trees: phylogenetic inference is complicated by incomplete lineage sorting and gene flow. *Mol Biol Evol.* 31:2004–2017.
- Kvikstad EM, Makova KD. 2010. The (r)evolution of SINE versus LINE distributions in primate genomes: sex chromosomes are important. *Genome Res.* 20:600–613.
- Lander ES, et al. 2001. Initial sequencing and analysis of the human genome. *Nature* 409:860–921.
- Li B, Zhang G, Willerslev E, Wang J. 2011. Genomic data from the Polar Bear (*Ursus maritimus*). *Gigascience*. Available from: <http://dx.doi.org/10.5524/100008>
- Li G, et al. 2013. Comparative analysis of mammalian Y chromosomes illuminates ancestral structure and lineage-specific evolution. *Genome Res.* 23:1486–1495.
- Li H, Durbin R. 2009. Fast and accurate short read alignment with Burrows-Wheeler transform. *Bioinformatics* 25:1754–1760.
- Li H, et al. 2009. The Sequence Alignment/Map format and SAMtools. *Bioinformatics* 25:2078–2079.
- Liu S, et al. 2014. Population genomics reveal recent speciation and rapid evolutionary adaptation in polar bears. *Cell* 157:785–794.
- McKenna A, et al. 2010. The Genome Analysis Toolkit: a MapReduce framework for analyzing next-generation DNA sequencing data. *Genome Res.* 20:1297–1303.
- Miller W, et al. 2012. Polar and brown bear genomes reveal ancient admixture and demographic footprints of past climate change. *Proc Natl Acad Sci U S A.* 109:E2382–E2390.

- Murphy WJ, et al. 2006. Novel gene acquisition on carnivore Y chromosomes. *PLoS Genet.* 2:e43.
- Nakagome S, Pecon-Slaterry J, Masuda R. 2008. Unequal rates of Y chromosome gene divergence during speciation of the family Ursidae. *Mol Biol Evol.* 25:1344–1356.
- O'Brien S, Menninger J, Nash W. 2006. *Atlas of mammalian chromosomes*. Hoboken (NJ): John Wiley & Sons.
- Pagès M, et al. 2008. Combined analysis of fourteen nuclear genes refines the Ursidae phylogeny. *Mol Phylogenet Evol.* 47:73–83.
- Pagès M, et al. 2009. A system for sex determination from degraded DNA: a useful tool for palaeogenetics and conservation genetics of ursids. *Conserv Genet.* 10:897–907.
- Posada D, Crandall K, Holmes EC. 2002. Recombination in evolutionary genomics. *Annu Rev Genet.* 36:75–97.
- Rogers J, Gibbs R. 2014. Comparative primate genomics: emerging patterns of genome content and dynamics. *Nat Rev Genet.* 15:347–359.
- Ross MT, et al. 2005. The DNA sequence of the human X chromosome. *Nature* 434:325–337.
- Rosser ZH, Balaesque P, Jobling M. 2009. Gene conversion between the X chromosome and the male-specific region of the Y chromosome at a translocation hotspot. *Am J Hum Genet.* 85:130–134.
- Sacks BN, et al. 2013. Y chromosome analysis of dingoes and southeast asian village dogs suggests a neolithic continental expansion from Southeast Asia followed by multiple Austronesian dispersals. *Mol Biol Evol.* 30:1103–1118.
- Scozzari R, et al. 2014. An unbiased resource of novel SNP markers provides a new chronology for the human Y chromosome and reveals a deep phylogenetic structure in Africa. *Genome Res.* 24:535–544.
- Skaletsky H, et al. 2003. The male-specific region of the human Y chromosome is a mosaic of discrete sequence classes. *Nature* 423:825–837.
- Slaterry JP, Sanner-wachter L, O'Brien SJ. 2000. Novel gene conversion between X-Y homologues located in the nonrecombining region of the Y chromosome in Felidae (Mammalia). *Proc Natl Acad Sci U S A.* 97:5307–5312.
- Smit AFA, Riggs AD. 1995. MIRs are classic, tRNA-derived SINES that amplified before the mammalian radiation. *Nucleic Acids Res.* 23:98–102.
- Smit AFA, Hubley R, Green P. 1996. RepeatMasker Open-3.0. 1996–2010 <<http://www.repeatmasker.org>>.
- Soh YQS, et al. 2014. Sequencing the mouse Y chromosome reveals convergent gene acquisition and amplification on both sex chromosomes. *Cell* 159:800–813.
- Trombetta B, Sellitto D, Scozzari R, Cruciani F. 2014. Inter- and intraspecies phylogenetic analyses reveal extensive X-Y gene conversion in the evolution of gametologous sequences of human sex chromosomes. *Mol Biol Evol.* 31:2108–2123.
- Van Oven M, Van Geystelen A, Kayser M, Decorte R, Larmuseau MHD. 2014. Seeing the wood for the trees: a minimal reference phylogeny for the human Y chromosome. *Hum Mutat.* 35:187–191.
- Veeramah KR, Hammer MF. 2014. The impact of whole-genome sequencing on the reconstruction of human population history. *Nat Rev Genet.* 15:149–162.
- Wallner B, et al. 2013. Identification of genetic variation on the horse Y chromosome and the tracing of male founder lineages in modern breeds. *PLoS One* 8:e60015.
- Walters-Conte KB, Johnson DLE, Allard MW, Pecon-Slaterry J. 2011. Carnivore-specific SINES (Can-SINES): distribution, evolution, and genomic impact. *J Hered.* 102(Suppl.):S2–S10.
- Wei W, et al. 2013. A calibrated human Y-chromosomal phylogeny based on resequencing. *Genome Res.* 23:388–395.
- Willard HF. 2003. Tales of the Y chromosome. *Nature* 423:810–813.
- Wilson AC, et al. 1985. Mitochondrial DNA and two perspectives on evolutionary genetics. *Biol J Linn Soc.* 26:375–400.

Associate editor: Kateryna Makova


Supplementary information

Genome-wide search identifies 1.9 megabases from the polar bear Y chromosome for evolutionary analyses


Tobias Bidon, Nancy Schreck, Frank Hailer , Maria A. Nilsson and Axel Janke

Details on *in-vitro* experiments

Polymerase chain reaction (PCRs) were performed in 15 µl reaction volumes containing 2x Taq DNA Polymerase mix (VWR International GmbH, BDH Prolabo, Darmstadt, Germany), 0.17 µg/µl BSA (New England Biolabs, Ipswich, MA, USA), 0.27 µM each of forward and reverse primer, and 10-15 ng template DNA. The amplification protocol started with 95°C for 3 min followed by 14 cycles at 94°C for 30 s, a touchdown step for 25 s (see Table S7 for specific starting temperatures) and 72°C for 75 s. This was followed by 25 cycles of 94°C for 30 s, a specific annealing temperature for 25 s (Table S7) and 72°C for 75 s. Final elongation was conducted for 10 min at 72°C.



A



B

Figure S1: Evidence for X/Y gene conversion events (yellow stars) in bears.

A Rooted phylogeny of ~397 bp from ZFY and ZFX sequences in bears and other mammals.
B Statistical haplotype network of bear ZFY and ZFX exon sequences (bear sequences from **A**).




Figure S2: Average-depth (AD) ratio for scaffolds in the polar bear assembly (enlarged version of Fig. 2A).



Figure S3: Annotations of Y-linked scaffolds.

Exons homologous to mouse and human are shown in red. Previously published Y-linked polar bear sequences are shown in pink. The repeat unit of each microsatellite is indicated and regions with >200 bp of consecutive "N" are highlighted in gray. Due to the high abundance, only placental mammalian non-LTR retrotransposons ≥500 bp (LINEs) and ≥100 bp (SINEs) were plotted. The maps of additional scaffolds are shown in **Figure 3**.

Table S1: Gene sequences used in the similarity search. Previously published polar bear sequences are in bold. Asterisk indicates the 18 mammalian genes that were found to have identity $\geq 80\%$ to scaffolds of the polar bear assembly (see Table 1).

No.	Gene	Sequence class in humans (according to Skaletsky et al. 2003)	Available sequences*	Fragment extracted	Species	GenBank accession number	Scaffold identified	Number of gene fragments recovered on respective scaffold
Y-linked								
1*	AMELY		Exon 5	exon	<i>Ursus maritimus</i>	AM941064.1	115	1 exon
	AMELY	X-degenerate	RefSeqGene	exon	<i>Homo sapiens</i>	NG_008011.1	389	2 exons
2	BPY2	Ampliconic	RefSeqGene	exon	<i>Homo sapiens</i>	NG_009862.1	-	-
3	CDY1	Ampliconic	RefSeqGene	exon	<i>Homo sapiens</i>	NG_011754.1	-	-
4*	DDX3Y (DBY)	X-degenerate	Intron 5 + partial CDS	exon	<i>Pan troglodytes</i>	JF293113.1	309	1 exon
	DDX3Y (DBY)		RefSeqGene	exon	<i>Homo sapiens</i>	NG_012831.1	184, 309	12 exons, 17 exons
	DDX3Y (DBY)		RefSeqGene	exon	<i>Mus musculus</i>	NM_012008.2	184, 309	9 exons, 15 exons
5*	EIF1AY	X-degenerate	mRNA	exon	<i>Homo sapiens</i>	NM_004681.2	20, 297	3 exons, 5 exons
	EIF1AY		Partial exon	exon	<i>Pan troglodytes</i>	AB176583.1	-	-
6*	EIF2S3Y		mRNA	exon	<i>Mus musculus</i>	NM_012011.1	20, 318, 646	8 exons, 6 exons, 2 exons
7	HSFY1	Ampliconic	RefSeqGene	exon	<i>Homo sapiens</i>	NG_012030.1	-	-
8*	KDM5D (SMCY)	X-degenerate	RefSeqGene	exon	<i>Homo sapiens</i>	NG_032920.1	297, 301, 318	12 exons, 9 exons, 8 exons
	KDM5D (SMCY)		mRNA	exon	<i>Mus musculus</i>	NM_011419.3	297, 301, 318	8 exons, 4 exons, 7 exons
	KDM5D (SMCY)		intron 4	complete (intron)	<i>Ursus maritimus</i>	AB261824.1	318	1 intron
9*	NLGN4Y	X-degenerate	RefSeqGene	exon	<i>Homo sapiens</i>	NG_028212.1	104	4 exons
10*	PCDH11Y	X-transposed	RefSeqGene	exon	<i>Homo sapiens</i>	NG_011652.1	186	3 exons
11*	PRKY	X-degenerate	mRNA	exon	<i>Homo sapiens</i>	Y15801.1	104	4 exons
12	PRY	Ampliconic	RefSeqGene	exon	<i>Homo sapiens</i>	NG_032916.1	-	-
13	RBM31Y		mRNA	exon	<i>Mus musculus</i>	NM_028970.1	-	-
14	RBM5Y		mRNA	exon	<i>Mus musculus</i>	NM_011253.2	-	-
15*	RBM5Y1A1	Ampliconic	RefSeqGene	exon	<i>Homo sapiens</i>	NG_012805.1	105, 369	1 exon, 6 exons
16*	RPS4Y		Partial CDS	exon	<i>Pan troglodytes</i>	AH012491.2	13, 253	1 exon, 6 exons
17*	RPS4Y2	X-degenerate	RefSeqGene	exon	<i>Homo sapiens</i>	NG_032924.1	253	5 exons
18*	SLY		mRNA	exon	<i>Mus musculus</i>	NM_201530.2	53	1 exon
19*	SRY		Exon 1	complete (exon +UTR)	<i>Ursus maritimus</i>	AM748305.1	579	1 exon + UTR
	SRY	X-degenerate	RefSeqGene	exon	<i>Homo sapiens</i>	NG_011751.1	-	-
	SRY		mRNA	exon	<i>Mus musculus</i>	NM_011564.1	-	-
20	SSTY1		mRNA	exon	<i>Mus musculus</i>	NM_009220.2	-	-
21	SSTY2		mRNA	exon	<i>Mus musculus</i>	NM_023546.3	-	-
22*	TBL1Y	X-degenerate	mRNA	exon	<i>Homo sapiens</i>	NM_033284.1	46, 104	1 exon, 12 exons
23	TGIF2LY	X-transposed	mRNA	exon	<i>Homo sapiens</i>	NM_139214.2	-	-
24	TMSB4Y	X-degenerate	mRNA	exon	<i>Homo sapiens</i>	NM_004202.2	-	-
25	TSPY1	Ampliconic	RefSeqGene	exon	<i>Homo sapiens</i>	NG_027958.1	-	-
26*	UBA1Y		mRNA	exon	<i>Mus musculus</i>	NM_011667.2	134, 403, 605, 4889	12 exons, 1 exon, 13 exons, 1 exon
	UBA1Y		Exon 18	complete (exon)	<i>Ursus maritimus</i>	AM748329.1	605	1 exon
27*	USP9Y	X-degenerate	RefSeqGene	exon	<i>Homo sapiens</i>	NG_008311.1	20, 184, 309, 318	1 exon, 34 exons, 38 exons, 1 exon
	USP9Y		mRNA	exon	<i>Mus musculus</i>	NM_148943.2	20, 184,	1 exon, 29 exons, 35

							309, 318	exons, 1 exon
28*	UTY		mRNA	exon	<i>Mus musculus</i>	NM_009484.2	134, 309	19 exons, 21 exons
29	VCY	Ampliconic	RefSeqGene	exon	<i>Homo sapiens</i>	NG_032915.1	-	-
30	XKRY	Ampliconic	RefSeqGene	exon	<i>Homo sapiens</i>	NG_032917.1	-	-
31*	ZFY		Final intron	complete (intron)	<i>Ursus maritimus</i>	AB261808.1	318	1 intron
	ZFY		partial gene	complete (partial gene)	<i>Ursus maritimus</i>	AM748297.1	20, 318	1 exon, 1 exon
	ZFY		Partial CDS	exon	<i>Canis lupus</i>	JX475923.1	20, 318	3 exons, 4 exons
	ZFY	X-degenerate	RefSeqGene	exon	<i>Homo sapiens</i>	NG_008113.1	20, 318	5 exons, 6 exons
32	ZFY1		mRNA	exon	<i>Mus musculus</i>	NM_009570.4	-	-
X-linked								
1	AMELX		Exon 5	exon	<i>Ursus maritimus</i>	AM941056.1	115, 6612	1 exon, 1 exon
2	ZFX		Final intron	complete (intron)	<i>Ursus maritimus</i>	AB261816.1	20	1 intron
3	ZFX		partial gene	complete (partial gene)	<i>Ursus maritimus</i>	AM941048.1	20, 318, 3838	1 exon, 1 exon, 1 exon

Table S2: Y-linked scaffolds <10 kb (n=92) identified by their AD-ratio.

Scaffold ID	Size (kb)	AD-ratio	Scaffold ID	Size (kb)	AD-Ratio
882	9	0	2560	1	0
949	6	0	2698	1	0
951	6	0	2697	1	0
955	6	0.085	2720	1	0
983	5	0	2812	1	0
1057	5	0	2853	1	0.136
1166	4	0	2856	1	0
1214	3	0	2882	1	0
1229	3	0.3	2879	1	0
1232	3	0	3005	1	0
1235	3	0	3008	1	0.225
1237	3	0	3026	1	0
1251	3	0	3027	1	0
1261	3	0	3043	1	0
1273	3	0	3094	1	0
1329	3	0	3105	1	0
1346	3	0	3158	1	0
1359	3	0	3258	1	0
1484	3	0	3283	1	0.299
1621	2	0	3349	1	0
1629	2	0	3381	1	0.271
1665	2	0	3481	1	0
1687	2	0	3487	1	0
1727	2	0	3578	1	0
1741	2	0	3585	1	0
1756	2	0	3598	1	0.196
1760	2	0	3608	1	0
1772	2	0	3649	1	0
1850	2	0	3662	1	0.163
1880	2	0	3691	1	0
1885	2	0	3693	1	0
1911	2	0	3723	1	0
1934	2	0	3838	1	0
1956	2	0	3849	1	0
2053	2	0	3886	1	0
2099	2	0	3889	1	0
2138	2	0	3894	1	0
2227	2	0	3958	1	0.184
2256	2	0	3965	1	0
2285	2	0	4014	1	0
2428	2	0	4021	1	0.182
2434	2	0	4107	1	0.172
2433	2	0	4138	1	0
2467	1	0	4147	1	0
2483	1	0	4146	1	0
2502	1	0	4157	1	0

Table S3: Scaffolds identified by the similarity search to be Y-linked, but with AD-ratios indicative of being autosomal or X-linked.

Scaffold ID	Size [Kbp]	AD-ratio	Inferred chromosomal location	Homology search
13	26,707	0.962	A	<i>RPS4Y</i>
20	22,125	1.942	X	<i>EIF1AY, EIF2S3Y, USP9Y, ZFY</i>
46	15,941	0.978	A	<i>TBL1Y</i>
53	14,458	0.973	A	<i>SLY</i>
104	6,801	0.969	A	<i>NLGN4Y, PRKY, TBL1Y</i>
105	6,717	1.944	X	<i>RBMV1A1</i>
115	5,608	1.934	X	<i>AMELY</i>
134	4,672	1.943	X	<i>UBA1Y, UTY</i>
184	2,589	1.949	X	<i>DDX3Y, USP9Y</i>
186	2,578	1.956	X	<i>PCDH11Y</i>
253	821	1.933	X	<i>RPS4Y, RPS4Y2</i>
301	351	1.928	X	<i>KDM5D</i>
4889 *	0.9	-	-	<i>UBA1Y</i>
6612 **	0.7	0	-	<i>AMELY</i>

* Reads mapped to scaffold 4889 with quality <20, so this AD-ratio could not be determined.

** Scaffold 6612 (749 bp) has an AD-ratio of zero but was filtered out due to its size <1kb.

Table S4: Scaffold combinations containing Y- and X-linked scaffolds

Details from the 10 query genes that were observed on ≥ 2 scaffolds are shown (see also Table S1).

Scaffold with AD-ratio indicative of Y-linkage and male-specific amplification in-vitro	Scaffold with putative X-linkage (AD-ratio in brackets)	Query Gene
297	20 (1.94)	<i>EIF1AY</i>
297	301 (1.94)	<i>KDM5D</i>
309	134 (1.94)	<i>UTY</i>
309	184 (1.95)	<i>DDX3Y</i>
309	20 (1.94), 184 (1.95)	<i>USP9Y</i>
318	20 (1.94)	<i>EIF2S3Y, ZFY</i>
318	301 (1.94)	<i>KDM5D</i>
318	20 (1.94), 184 (1.95)	<i>USP9Y</i>
369	105 (1.94)	<i>RBM1A1</i>
389	115 (1.93)	<i>AMELY</i>
403	134 (1.94)	<i>UBA1Y</i>
605	134 (1.94)	<i>UBA1Y</i>
646	20 (1.94)	<i>EIF2S3Y</i>
6612	115 (1.93)	<i>AMELY</i>

Table S5: Percentage of transposable elements in 1.6 Mb Y-chromosomal scaffolds in comparison to the genome-wide average in polar and brown bears.

	Polar bear		Brown bear
	Y-scaffolds	Genome	Genome
SINE	8.23	8.18	8.15
MIRs	0.42	3.03	3.05
CanSINE	7.81	5.05	5.10
LINE	38.88	21.03	21.09
LINE-1	38.42	16.93	16.96
LINE-2	0.44	3.58	3.60
LINE-3	0.03	0.39	0.39
RTE	0.00	0.12	0.13
ERV	5.29	5.40	5.45
DNA transposons	1.98	3.00	3.03
Total	54.38	39.20	39.27

Table S6: Sequence characteristics of microsatellites with ≥ 15 repeat units on Y-linked scaffolds. Bases in lowercase denote regions with imperfect tandem repeats.

[illegible]

[illegible]

scaffold389	TG	20	12691	TGTGTGTGtcTGTGTGTGTGtaTGTGTGTGTGTGTGTGTGTG
scaffold389	TG	20	22172	TGTGTGTGTGTGTGTGTGTGTGTGTGTGTGTGTGTGTGT
scaffold389	TG	21	26235	TGTGTGTGTGTGTGTGTGTGTGTGTGTGTGTGTGTGTGT
scaffold389	TG	20	52851	TGTGTGTGTGTGTGTGTGTGTGTGTGTGTGTGTGTGTGT
scaffold389	AC	17	73174	ACACACACACACACACACACACACACACACACACACAC
scaffold389	TG	18	74158	TGTGTGTGgcTGTGTGTGTGTGTGTGTGTGTGTGTGT
scaffold389	TG	19	76543	TGTGTGtagTGTGTGTGTGTGTGTGTGTGTGTGTGTGTG
scaffold393	AC	20	1822	ACACACACACACACACACACACACACACACACACACAC
scaffold393	GT	15	13861	GTGTGTGTGTGTGTGTGTGTGTGTGTGTGTGTGTGT
scaffold393	CT	16	22325	CTCTCTCTCTCTCTCTCTCTCTCTCTCTCTCTc
scaffold393	GT	18	48001	GTGTGTGTGTGTGTGTGTGTGTGTGTGTGTGTGTGTG
scaffold393	CT	14	54007	CTCTtCTCTCTCTCTCTCTCTCTCTCTCTCTCT
scaffold393	CA	24	54039	CACACACACACACACACACACACACACACACACAtaCACACACACA CACA
scaffold393	GT	16	61378	GTGTGTGTGTGTGTGTGTGTGTGTGTGTGTGTGTG
scaffold403	GT	17	16279	GTGTGTGTGTGTGTGTGTGTGTGTGTGTGTGTGTG
scaffold403	GT	19	26610	GTGTGTGTGTGTGTGTGTGTGTGTGTGTGTGTGTGTGT
scaffold403	AC	15	44337	ACACACACACACACACACACACACACACACACa
scaffold403	AC	15	51140	ACACACACACACACACACACACACACACACACa
scaffold420	TG	16	7834	TGTGTGTGTGTGTGTGTGTGTGTGTGTGTGTGTG
scaffold420	CAA	15	29461	CAACAACAACAACAACAACAACAACAACAACAACAACA
scaffold519	AAGG	23	1420	AAGGAAGGAAGGAAGGAAGGAAGGAAGGAAGGAAGGAAG GAAGGAAGGAAGGAAGGAAGGAAGGAAGGAAGGAAGGaaaaagAAGG AAGGAAGGAAGG
scaffold579	T	26	10424	TTgTTTTTTTTTTTTTTTTTTTTTTTTTTTTTTTT
scaffold579	T	33	11001	TTTTTTTTTTTTTTTTTTTTTTTTTTTTTTTTTTTT
scaffold579	AAGA	35	13351	AAGAaggaAAGAaggaAAGAAAGAAAGAAAGAAaAAGAAAAAAGAAAGAAAG AagaAAGAAAGAAAGAAAGAAAGAAAGAAAGAAAGAAAGAAgaAAGAAAG AAAGAAAGAAAGAAAGAAAGAAAGAAAGAAAGAAAGAAAGAAAGAAAGAAAGAA AAGAAAGAAAGAtaaAAGAAAGAaa
scaffold579	A	17	14764	AAAAAAAAAAAAAAAAAAAA
scaffold579	TC	17	16925	TCTCTCTCTCTCTCTCTCTCTCTCTCTCTCTCTC
scaffold657	AC	24	6191	ACACaaACACACACACACACACACACACACACACACACACACA CACACa

Table S7. Primer sequences and PCR conditions of fragments used for *in-vitro* validation of Y-scaffolds.

Scaffold	Forward primer (5'→3')	Reverse primer (5'→3')	Fragment size (bp) incl. primers	T _{A1} (°C) 14 cycles	T _{A2} (°C) 25 cycles
297	ACAGAGAGATGCGGTCCTG	GGAGACTCTGCTGACCAAGG	783	69	62
309	TCAGCCTTGCTTCATTGTTTC	ATTGTGTGCCTGTTCCATTC	696	68	61
318.20	CAATGCTAGAGATGCAAGTGG	AAGCATATGAAGACACACAGTGG	757	67	60
318.21	ATGCTATGATCCAGCAATCC	TCCTTGTTAATTCTTCTCTGATGC	800	67	60
318.22	CTGACATACGTGGCTCACAG	ATGTGGAGGTTCAAGGAGGAC	800	67	60
318.23	GTGGTCAGATCCAGCTCCTC	GACCTTGCTTCTGCCTTCTC	786	67	60
322**	GAGTAGAGCTGGTGCTTGTGAG	GAAGCAGAGCTCAAGTCTGAAG	687	68	61
369	CATTGAACAAGGGCACATC	TCCTCTAGGAACCCAACTGC	780	68	61
393	AATTGTGAAGGATGAGGAAGG	CCAGAGAAGCAGAGGATGG	788	70	63
389**	ACCCACTGCTGTTCTGTATCC	CCAACAGTGTAGTGGTTGTGC	679	68	61
403**	CACTCAGGAGAGCACAGGTC	TGTGTGTCGTAAGCAGAGGTC	796	68	61
420	TTCAGAGGGAAGGGTGTAGG	AATGGTAGTGAGGAGGTATGG	635	69	62
519	CCCAAGTTAGGGAAGTTTGC	TGTTTCAGCTTCAGCTGTCC	734	68	61

579*	CTGCAGGCCTGTCAATGTTA	TGTGTATCGACCCATACTTTG	660	66	61
605	TTTGACGGCTGAGCAATATC	GATGCAGCCACAATGAAATC	706	68	61
646	TGTCTAGCCATCTGGTCCTG	CATGGACATTGTTGCATTGA	778	69	62
657	TCCAGGTTGTCAAGCACATG	CTTGCTCCATCCACATGCTG	769	69	62
596	ATCACCTCTCCACTCACAA	CCACCTCTTTGACTTCCTGG	735	69	62
613	TTGTGGACATTGCTGCTTCT	TCTGAACATAGGCTGCAACC	674	69	62
632	CTATCGTGGACATTGCTGCT	CTATCGTGGACATTGCTGCT	621	69	62
771	AGAACTTCCTGTACCTGGCT	GCTGATTCCGTGGATGTTCA	753	69	62
795	GGAACACGACAAGGATCTCC	TTGTGCCAATACCATGCTGT	705	69	62
813	ATCTTGCTGCTCCTCTGTTG	CCACGTCACATCATACTGCA	767	69	62
236a	GACTCAAGCTCAGCGTTCAC	GTCCAGTGAAGAGGTAGGC	760	68	61
267a	ATACATACGCGCACACACAC	GCACATACCTCGTCAAGGAC	655	68	61
253x	CTGGAGGTCTGTGTGTGGAG	AACCTAATGCGTGTGTCTGC	796	68	61
301x	TGGATAAGGCATTCTGCAAG	CTGAAGGCAGTGGTGAAGAG	717	68	61

* Primers from (Bidon et al. 2014) (fragment 579.1B)

** Primers from (Kutschera et al. 2014)

a Autosomal scaffold based on AD-ratio

x X-linked scaffold based on AD-ratio

Table S8. Accession numbers and sample origin of polar, brown and black bear genomes

Species	Sample origin	Number of individuals	Accession numbers
<i>Ursus maritimus</i>	Spitsbergen, Svalbard	9	SRX155945, SRX155949, SRX155951, SRX155953, SRX155954, SRX155955, SRX155957, SRX155960, SRX155961
	Alaska	3	SRX156102, SRX156103, SRX156105
<i>Ursus arctos</i>	ABC-Islands	1	SRX156108
<i>Ursus americanus</i>	Alaska	1	SRX156137